

# Phylogeny and phylogeography of Atlantic oyster species: evolutionary history, limited genetic connectivity and isolation by distance

C. Lazoski<sup>1</sup>, J. Gusmão<sup>1,2</sup>, P. Boudry<sup>3</sup>, A. M. Solé-Cava<sup>1,\*</sup>

<sup>1</sup>Laboratório de Biodiversidade Molecular – Departamento de Genética, Instituto de Biologia, UFRJ, CCS, Bloco A, sala A2-98, Ilha do Fundão, 21941-490 Rio de Janeiro, Brazil

<sup>2</sup>Laboratório de Genética Marinha — Departamento de Genética, Instituto de Biologia Roberto Alcântara Gomes, UERJ, Maracanã, 20550-900 Rio de Janeiro, Brazil

<sup>3</sup>Iremer, UMR M100 Physiologie et Ecophysiologie des Mollusques Marins, 29280 Plouzané, France

**ABSTRACT:** The phylogenetic relationships between naturally occurring Atlantic *Crassostrea* oyster species were inferred through analyses of mitochondrial (cytochrome oxidase subunit I and 16S) and nuclear (second internal transcribed spacer) sequences. We also scored 15 allozyme loci on 422 oysters to study population structuring of *C. rhizophorae* and *C. brasiliana* along 9000 km of the Western Atlantic coastline. Despite morphological similarities, *C. virginica* was genetically more closely related to *C. rhizophorae* than to *C. brasiliana*. In contrast, *C. paraibanensis* was genetically indistinguishable from *C. brasiliana*, which is probably a junior synonym of the African *C. gasar*. Significant genetic differentiation between populations of *C. rhizophorae* and *C. gasar* were found along the Western Atlantic coast, supporting an isolation-by-distance pattern.

**KEY WORDS:** Population genetics · Biogeography · Allozymes · Cytochrome *c* oxidase subunit I · COI · 16S · Internal transcribed spacer 2 · ITS-2

Resale or republication not permitted without written consent of the publisher

## INTRODUCTION

Oysters are the world's most economically important mollusks, with a global average production of 4.6 million tons yr<sup>-1</sup> (FAO 2010). In the Western Atlantic, oysters of the genus *Crassostrea* are the fourth largest marine fishery resource (average of 200 000 tons yr<sup>-1</sup>; FAO 2010). The most heavily exploited species in the region are *C. virginica* (Gmelin, 1791), *C. rhizophorae* (Guilding, 1828) and the exotic *C. gigas* (Thünberg, 1793).

Phylogenetic studies have demonstrated the existence of 2 evolutionary lineages within the *Crassostrea* genus, one from the Atlantic and another from the Pacific, which could have been formed during the closure of the Tethys Seaway (Ó Foighil & Taylor 2000). Molecular phylogenies of both nuclear and mitochondrial genomes have helped to clarify the taxonomic

confusion of the Pacific *Crassostrea* species (Reece et al. 2008, Wu et al. 2010). On the other hand, the taxonomic status of Atlantic *Crassostrea* oysters is still confused: there is clear evidence for the existence of at least 2 oyster species along the Brazilian coast (Absher 1989, Ignacio et al. 2000), but 5 other nominal species have been cited for the area, regarded as native (*C. virginica*, Carriker & Gaffney 1996; *C. paraibanensis*, Singarajah 1980) or invasive (*C. gigas*, Melo et al. 2010b; *C. gasar* Adanson, 1757, Lapègue et al. 2002; *Crassostrea* sp., Varela et al. 2007, Melo et al. 2010a) species. The correct identification of those oyster species is of paramount importance, because they are used in aquaculture and in biomonitoring studies (e.g. Rebelo et al. 2003), and may be under environmental threat (Carranza et al. 2009).

The uncertainty in the classification of Atlantic oyster species may have been caused in part by the great

\*Corresponding author. Email: sole@biologia.ufrj.br

morphological and ecological similarities between *Crassostrea brasiliana* (Lamarck, 1819) and *C. virginica*. *C. rhizophorae* and *C. brasiliana* are also morphologically quite similar and have been considered synonyms in monographic studies of Brazilian mollusks (Rios 1994). This may have led some systematists to suggest that *C. rhizophorae* and *C. brasiliana* were morphotypes of a single species (*C. virginica*) (Harry 1985, Carriker & Gaffney 1996).

*Crassostrea rhizophorae* is found in the intertidal zone, either attached to *Rhizophora mangle* roots or on rocks, from the Caribbean to the southern Atlantic (Santa Catarina State, Brazil) (Carriker & Gaffney 1996). *C. brasiliana* usually attaches to rocks in infralittoral zones, and it is most abundant from southeastern Brazil (Espírito Santo State) to the more temperate regions in southern Brazil (Absher 1989). The species has also been observed in French Guyana (Lapègue et al. 2002) and in Venezuela (Hoover & Gaffney 2005).

Besides the ecological differences, these oyster species can be distinguished by shell size, which is much larger in *Crassostrea brasiliana* (50 to 190 mm) than in *C. rhizophorae* (20 to 65 mm) (Absher 1989). However, this only holds true for adult specimens. Based on Lamarck's small-shelled holotype for *C. brasiliana*, Singarajah (1980) considered it a synonym of *C. rhizophorae*. Hence, he described the large oysters found in Paraíba State (northeastern Brazil) as a new species, *C. paraibanensis* (Singarajah 1980). However, other authors considered that this species might be synonymous with *C. brasiliana* (Rios 1994, Carriker & Gaffney 1996).

Those taxonomic problems can also result from differing views on the degree of differentiation required to define a species in *Crassostrea*. Since the late 1970s, many studies have employed molecular methods to address the systematics and population genetics of *Crassostrea* species. For example, consistent genetic differences have been found between *C. rhizophorae* and *C. virginica* using allozyme (Buroker et al. 1979, Hedgecock & Okazaki 1984) and ribosomal DNA data (Littlewood 1994), as well as between *C. rhizophorae* and *C. brasiliana* using morphology (Absher 1989), allozymes (Ignacio et al. 2000, Lazoski 2004) and mitochondrial 16S or cytochrome *c* oxidase subunit I (COI) data (Lapègue et al. 2002, Boudry et al. 2003, Lazoski 2004, Varela et al. 2007). In contrast, Lapègue et al. (2002) did not find any differences between 16S sequences of West African *C. gasar* and South American *Crassostrea* specimens, which led them to suggest that *C. brasiliana* and *C. gasar* might be the same species (with *C. brasiliana* being a junior synonym of *C. gasar*). However, no molecular studies have compared *C. virginica* and *C. brasiliana* populations.

Although marine species with potentially high dispersal capability are believed to have high levels of

gene flow, there are many examples showing the opposite (e.g. Palumbi 2003, Johansson et al. 2008), including studies of *Crassostrea* species (Hedgecock & Okazaki 1984, Hare & Avise 1998, Xiao et al. 2010). Population genetic studies have unveiled patterns of isolation by distance among oyster populations (Launey et al. 2002, Rose et al. 2006, Xiao et al. 2010), and are of great interest not only when addressing evolutionary and ecological processes but also as a basis for management and conservation of commercially important marine species.

In spite of the economic importance of *Crassostrea* oysters, no study has been carried out on the degree of genetic connectivity of their populations along the Western Atlantic. Moreover, the unresolved systematics of those species has confounded the compiled fishery statistics data (FAO 2010) for the Latin American coast, and the true geographic distribution of oyster species along this area is still uncertain (Singarajah 1980, Carriker & Gaffney 1996, Ignacio et al. 2000, Lapègue et al. 2002, Varela et al. 2007).

In this paper, we compare Atlantic populations of *Crassostrea* by means of allozyme electrophoresis and analyses of nuclear (internal transcribed spacer 2, ITS-2) and mitochondrial (COI, 16S) sequences. Our objectives were to define the taxonomic boundaries among the Western Atlantic oysters, *C. brasiliana*, *C. virginica*, *C. rhizophorae* and *C. paraibanensis*, and to verify their actual distribution. We also compared the nuclear and mitochondrial gene sequences from specimens of *C. brasiliana* and the African *C. gasar* to resolve their taxonomic ambiguity. Additionally, we investigated the levels of genetic variation and the population structures in *C. brasiliana* and *C. rhizophorae* along 9000 km of the Western Atlantic coastline.

## MATERIALS AND METHODS

**Sampling.** A total of 419 individuals, presumed to be either *Crassostrea rhizophorae* or *C. brasiliana*, were collected between May 1996 and February 2003 from 21 localities in Brazil and Panama (Table 1). Sixty-five samples of other *Crassostrea* species were also collected in the Gulf of Mexico (USA; 29° 03' N; 95° 07' W: *C. virginica*, N = 6), Africa (Senegal; 14° 10' N; 16° 51' W: *C. gasar*, N = 7; Lapègue et al. 2002) and Brazil (Itajaí; 26° 55' S; 48° 38' W: hatchery of *C. gigas*, N = 31; and Manguape, PB1, 07° 06' S; 34° 54' W: *C. paraibanensis*, N = 21) (Table 1, Fig. 1). Twenty-two individuals of *Ostrea* sp. from Brazil (Sepetiba, RJ1, 22° 58' S; 43° 42' W; Picinguaba, SP1, 23° 22' S; 44° 50' W) were used as an outgroup for the allozyme analysis. After collection, the oysters were transported (alive, in liquid N<sub>2</sub> or in ethanol) to the laboratory, where they were identified morpholog-

Table 1. *Crassostrea rhizophorae* and *C. brasiliiana*. Sampling sites, coordinates and habitats for the putative *Crassostrea rhizophorae* and *C. brasiliiana* collected. (i): intertidal zone; (s): subtidal zone. \*Samples available only for DNA analysis

Sites	Coordinates	Habitat	<i>C. brasiliiana</i>	<i>C. rhizophorae</i>
Panama				
Panama (PAN)	09°10'N; 80°17'W	Rocks (i)	–	14
Brazil				
Belém (PA)	01°25'S; 48°28'W	Mangrove (i)	12	–
Araioses (MA)	02°53'S; 41°54'W	Mangrove (i)	12*	–
Parnaíba (PI)	02°51'S; 41°45'W	Rocks (i)	5*	–
Fortaleza (CE)	03°46'S; 38°26'W	Rocks (i)	–	12*
Natal (RN1)	05°45'S; 35°11'W	Mangrove (i)	–	35
Natal (RN2)	05°52'S; 35°09'W	Rocks (i)	–	20
Mamanguape (PB1)	07°06'S; 34°54'W	Mangrove (i)	7	10
Cabo Branco (PB2)	08°07'S; 34°52'W	Mangrove (i)	–	24
Recife (PE)	08°46'S; 34°56'W	Mangrove (i)	24(1*)	1(3*)
Salvador (BA1)	13°00'S; 38°26'W	Rocks (i)	20	–
Caravelas (BA2)	17°44'S; 39°15'W	Mangrove (i)	–	30
Vitória (ES)	20°18'S; 40°17'W	Rocks (i)	–	12*
Sepetiba (RJ1)	22°58'S; 43°42'W	Rocks (i)	–	22
Guaratiba (RJ2)	22°58'S; 43°40'W	Mangrove (i)	–	20*
Rio das Ostras (RJ3)	22°31'S; 41°56'W	Rocks (s)	5*	–
Picinguaba (SP1)	23°22'S; 44°50'W	Rocks (i)	–	10
Itanhaém (SP2)	24°11'S; 46°47'W	Mangrove (i)	4	–
Cananéia (SP3)	25°00'S; 47°56'W	Rocks (s)	7*	–
Paranaguá (PR1)	25°32'S; 48°22'W	Rocks (s)	40	–
Paranaguá (PR1)	25°32'S; 48°22'W	Rocks (i)	–	35
Guaratuba (PR2)	25°51'S; 48°35'W	Rocks (i)	2	32
<b>Total</b>			<b>139</b>	<b>280</b>

ically (Absher 1989; Fig. 2), measured (shell height, length and depth), dissected, and stored in either liquid nitrogen or ethanol until required for genetic analyses.

**Allozymes.** Horizontal 12.5% starch gel electrophoresis was carried out as previously described (Ignacio et al. 2000). The 11 enzyme systems investigated and the 3 buffer systems used are summarized in Table 2. The enzyme systems were stained according to standard procedures (Manchenko 1994) and corresponded to a total of 15 allozyme loci.

**DNA purification, amplification and sequencing.** Total DNA purification was performed using a modified 1% N-cetyl N,N,N-trimethylammonium bromide (CTAB) protocol (2% CTAB, 20 mM EDTA, 0.2%  $\beta$ -mercaptoethanol, 0.1 M Tris, 1.4 mM NaCl), followed by a sodium acetate- and isopropanol-induced precipitation step (Gusmão & Solé-Cava 2002). Polymerase chain reactions (PCR) used approximately 10 ng of template DNA, 1 unit of *Taq* polymerase (GE Life Sciences), 200  $\mu$ M each of the 4 dinucleo-



Fig. 1. *Crassostrea* spp. and *Ostrea* sp. Collection sites for specimens




Fig. 2. *Crassostrea* spp. and *Ostrea* sp. Shells from Western Atlantic populations. *C. brasiliensis* (Brazil): **1** (SC); **2, 17** (PR1); **3** (PR2); **9, 18** (PE); **11** (PA); *C. rhizophorae* (Brazil): **4, 5** (RJ1); **6, 8** (RN2); **7** (SP1); **10, 20** (PR1); *C. virginica* (USA): **19** (Gulf of Mexico); *C. gigas* (Brazil): **16** (SC); *Ostrea* sp. (Brazil): **12, 13** (RJ1); **14, 15** (SP1). Unit scale: cm

Table 2. Enzymes studied (and abbreviations), Enzyme Commission numbers (EC), and buffer systems analyzed. TEM: 0.10 M Tris, 0.01 M EDTA, 0.10 M maleate, pH 7.4; TC8: 0.25 M Tris, 0.06 M citrate, pH 8.0; LI: 0.005 M citrate, 0.03 M Tris (gel), 0.06 M LiOH, 0.30 M borate (buffer tank), pH 8.5/8.1 (Manchenko 1994)

Enzymes	Abbrev	EC	Buffers
Adenylate kinase	AK	2.7.4.3	TEM
Catalase	CAT	1.11.1.6	TC8
Phosphoglucomutase	PGM	2.7.5.1	TC8
Phosphogluconate dehydrogenase	PGD	1.1.4.4	TEM
Phosphoglucose isomerase	PGI	5.3.1.9	TC8
Glutamate oxaloacetate transaminase	GOT	2.6.1.1	LI
Isocitrate dehydrogenase	IDH	1.1.1.42	TEM
Leucine amino peptidase	LAP	3.4.1.1	LI
Malate dehydrogenase	MDH	1.1.1.37	TC8
Mannose phosphate isomerase	MPI	5.3.1.8	TC8
Proline-phenylalanine dipeptidase	PEP-A	3.4.13.18	TC8

tides, 500 nM of each primer, and 1.5 mM MgCl<sub>2</sub> in 20 µl of 1× PCR buffer (GE Life Sciences). Amplifications were performed in a mini-cycler (Sprint) programmed to begin with a denaturing step of 3 min at 95°C, which was followed by 30 cycles consisting of

the following steps: 94°C for 1 min, 52°C for 1 min, and 72°C for 1 min, and a final extension step at 72°C for 5 min. Negative controls, involving template-free reactions, were included in all PCR amplifications. The primers used for the amplification of each specific genomic region were (1) 16SAR (5'-CGCCTGTTTATCAAAAACAT-3') and 16SBR (5'-CCG GTC TGA ACTCAG ATC ACG T-3') (Palumbi 1996) for the amplification of a 560 bp fragment of the mitochondrial large ribosomal subunit (16S); (2) PH19 (5'-CATCGACAC TT(T/C)GAA CGC A-3') and ITS2 (5'-AAT CCTGGTTAG TTT CTT TTC CTC CGCT-3') (Dixon et al. 1995) for the amplification of about 650 bp of the ITS-2; (3) LCO (5'-GGTCAA CAA ATC ATA AAG ATATTGG-3') and HCO (5'-TAA ACT TCA GGGTGA CCA AAAAAT CA-3') (Folmer et al. 1994) for the amplification of a 700 bp fragment of the COI mitochondrial gene; and (4)

COIbrF (5'-GGG TTT TGAGCA GTT TTA GCCGGG-3') and COIbrR (5'-GGT CAT CCA GAA GTGTAC GTC C-3'), developed in this work, for the *C. brasiliensis*-specific amplification of a 661 bp fragment of COI (annealing temperature of 60°C).

DNA sequencing was carried out as previously described (Gusmão et al. 2006). The purification of PCR products was performed with a GFX™ PCR DNA and Gel Band Purification Kit (GE Life Sciences) according to the manufacturer's instructions. We directly sequenced 47 ITS-2, 167 COI, and 24 16S DNA fragments from *Crassostrea* species originating in 24 populations. Sequences were deposited in GenBank (accession numbers FJ544267–FJ544312, FJ717606–FJ717651). Additionally, *Crassostrea* sequences from GenBank were included in our phylogenetic analyses (accession numbers EU007484, EU007485, EU007509, EU007511, Reece et al. 2008; NC\_007175, Milbury & Gaffney 2005; AJ312937, Lapègue et al. 2002; DQ839413, DQ839414, DQ839415, Pie et al. 2006). Sequences were aligned using the Clustal X multiple alignment program version 1.83 (Thompson et al. 1997), and alignments were confirmed by visual inspection.

**DNA sequence analysis.** Phylogenetic analyses were conducted using MEGA 4.0 (Tamura et al. 2007) and PAUP 4.0 (Swofford 1998) programs. Pairwise Kimura 2-parameter distances (K2P; Kimura 1980) were used to build neighbor-joining trees (Saitou & Nei 1987). The MODELTEST 3.06 program (Posada & Crandall 1998) was used to evaluate the most appropriate model of DNA substitution for maximum-likelihood analyses (Felsenstein 1981) of the data set. The best-fit models, which were chosen after comparisons between likelihood scores from different DNA substitution models, were the HKY model (HKY+G: A = 0.2233; C = 0.1764; G = 0.1993; T = 0.4010; gamma = 0.1600) for the COI analyses, and the K80 model (K80+G: A = 0.2233; C = 0.1764; G = 0.1993; T = 0.4010; gamma = 0.7191) for the ITS-2 analyses. Starting tree(s) were obtained via neighbor-joining, from which a heuristic search was employed using the branch-swapping algorithm (TBR, tree-bisection-reconnection). Branch support was assessed by bootstrapping the original data set using 1000 replicates.

**Divergence times, population subdivision, isolation by distance and historical demography.** Allozyme genotype frequencies were used to estimate allele frequencies, level of genetic variation (heterozygosity,  $H$ ), fits to Hardy-Weinberg equilibrium ( $F_{IS}$ ; Wright 1978), inbreeding indices ( $F_{ST}$ ; Wright 1978), and pairwise unbiased genetic distances ( $D_{Nei}$ ; Nei 1978) using the programs BIOSYS-1 version 1.7 (Swofford & Selander 1981) and GENETIX version 4.05 (Belkhir et al. 2002). The significance of  $F_{IS}$  ( $H_0: F_{IS} = 0$ ) was estimated using a  $\chi^2$  test (Waples 1987). Divergence times

were estimated from allozyme data using the relationship  $t = 5D_{Nei} \times 10^6$  (Nei 1987).

Spatial analysis of molecular variance (SAMOVA; Dupanloup et al. 2002) was conducted with both allozyme and COI markers to determine hierarchical genetic structures. The significance of the fixation indices against the null hypothesis of panmixis was also evaluated using a Markov-Chain procedure with 10 000 permutations, using the software ARLEQUIN version 3.11 (Excoffier et al. 2005). To test for nonrandom associations between genetic and geographic distances in *Crassostrea rhizophorae* and *C. brasiliensis* populations, we used the web service IBDWS version 3.11 (Jensen et al. 2005, <http://ibdws.sdsu.edu>), testing the significance of pairwise correlations between  $\Phi_{ST}$  (or  $F_{ST}$ , for allozyme data) and geographic distances (in km) between all populations, through a Mantel test (10 000 permutations). Geographic coordinates and distances between sampling sites, measured as the shortest spherical distance by sea, were obtained using Google Earth.

Haplotype ( $h$ ) and nucleotide ( $\pi$ ) diversity estimates, and 2 tests of neutrality, Tajima's  $D$  (Tajima 1989) and Fu's  $F_S$  (Fu 1997), were obtained from ARLEQUIN. For the COI data, we constructed a haplotype network using the statistical parsimony procedure of Templeton et al. (1992) implemented in the TCS version 1.21 program (Clement et al. 2000). The time of divergence between species was estimated, for mitochondrial sequences, using an evolutionary rate of 1.21% sequence divergence (K2P) per million years for COI (Marko 2002).

We used mismatch distribution analysis (Rogers & Harpending 1992), in the ARLEQUIN software, to test for historical population expansion events within oyster populations. If the sudden expansion model was not rejected, then the parameter tau ( $\tau$ ) was converted to time since expansion ( $t = \tau/2\mu$ ) in years before present (for 665 bp,  $\mu = 8.05 \times 10^{-6}$  substitutions per locus per year, assuming a 1 yr generation time and a 1.21% substitution rate; Marko 2002).

We used the coalescent-based program MDIV (Nielsen & Wakeley 2001), on CBSU Web Computing Resources (<http://cbsuapps.tc.cornell.edu/mdiv.aspx>), to estimate the maximum likelihood values of theta ( $\theta = 2N_{ef}\mu$ ), scaled migration rate ( $M = N_{ef}m$ ), scaled time of divergence ( $T = t_1/N_{ef}$ ), and time to most recent common ancestor (TMRCA =  $t_2/N_{ef}$ ) ( $N_{ef}$  = effective female population size,  $m$  = migration rate,  $t_1$  = population divergence time,  $t_2$  = gene coalescence time, and  $\mu$  = mutation rate). The coalescent-scaled parameter  $T$  was converted to  $T_{div}$  (time in years since 2 populations diverged) according to the formula:  $T_{div} = T\theta/(2\mu)$ . For each pairwise comparison, a minimum of 3 chains were run using the finite



Table 3 (continued)

Locus	<i>C. rhizophorae</i>										<i>C. brasiliana</i>					<i>C.p.</i> PB1	<i>C.v.</i> USA	<i>C.g.</i> SC	<i>Ostrea sp.</i>		
	PAN	RN1	RN2	PB1	PB2	BA2	RJ1	SP1	PR1	PR2	PA	PB1	PE	BA1	SP2				PR1	RJ1	SP1
<b>Lap-2</b>																					
(N)	(14)	(15)	(20)	(10)	(23)	(27)	(22)	(10)	(19)	(32)	(12)	(7)	(24)	(20)	(4)	(34)	(21)	(6)	(31)	(12)	(10)
A	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	-	-	-
B	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.00	-	-
C	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.00	1.00
<b>Mdh-1</b>																					
(N)	(14)	(15)	(20)	(10)	(23)	(27)	(22)	(10)	(19)	(32)	(12)	(7)	(24)	(20)	(4)	(34)	(21)	(6)	(31)	(10)	(10)
A	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.00	-	-	-
B	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.30	0.30
C	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	-	-	0.70	0.65
D	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.00	-	0.05
<b>Mdh-2</b>																					
(N)	(14)	(15)	(20)	(10)	(23)	(27)	(22)	(10)	(19)	(32)	(12)	(7)	(24)	(20)	(4)	(34)	(21)	(6)	(31)	(10)	(10)
A	0.25	0.03	0.02	-	0.02	0.06	0.09	-	-	0.06	0.08	0.14	0.17	0.15	-	0.01	0.14	-	-	-	-
B	-	-	-	-	-	-	-	-	0.03	-	-	-	-	-	-	0.03	-	-	-	-	-
C	0.75	0.97	0.98	1.00	0.98	0.94	0.91	1.00	0.97	0.94	0.92	0.86	0.83	0.85	1.00	0.96	0.86	1.00	1.00	-	-
D	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.75	1.00
E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.25	-
<b>Mpi</b>																					
(N)	(13)	(15)	(20)	(9)	(21)	(25)	(22)	(10)	(19)	(32)	(12)	(7)	(23)	(20)	(4)	(34)	(21)	(6)	(31)	(12)	(10)
A	-	-	-	-	-	-	-	-	-	-	-	-	0.02	0.03	-	-	-	-	0.31	-	0.05
B	0.04	-	-	-	-	-	-	-	-	-	0.33	0.50	0.46	0.52	0.13	0.13	0.29	-	0.68	1.00	0.95
C	0.15	0.10	-	-	0.07	0.20	0.11	-	0.18	0.09	0.59	0.43	0.48	0.45	0.87	0.84	0.62	0.33	0.01	-	-
D	0.27	0.67	0.55	0.56	0.50	0.40	0.46	0.35	0.58	0.81	0.08	0.07	0.04	-	-	0.03	0.09	0.33	-	-	-
E	0.50	0.23	0.40	0.39	0.38	0.24	0.41	0.55	0.19	0.10	-	-	-	-	-	-	-	0.33	-	-	-
F	0.04	-	0.05	0.05	0.05	0.12	0.02	0.10	0.05	-	-	-	-	-	-	-	-	-	-	-	-
G	-	-	-	-	-	0.04	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<b>Pep-1</b>																					
(N)	(14)	(15)	(20)	(10)	(23)	(25)	(21)	(6)	(19)	(32)	(12)	(7)	(24)	(20)	(4)	(34)	(21)	(6)	(31)	(12)	(3)
A	0.03	-	-	-	-	0.12	-	-	-	-	-	-	-	-	-	-	-	-	-	0.08	-
B	0.18	0.33	0.05	0.25	0.28	0.12	0.45	0.25	0.60	0.33	-	-	-	-	-	-	-	-	0.29	0.25	0.50
C	0.79	0.60	0.92	0.50	0.59	0.48	0.55	0.75	0.40	0.67	-	-	-	-	-	-	-	-	0.32	0.13	-
D	-	0.07	0.03	0.25	0.11	0.26	-	-	-	-	-	0.07	0.02	-	-	-	0.02	1.00	-	0.17	-
E	-	-	-	-	-	0.02	-	-	-	-	0.83	0.50	0.48	0.30	0.25	0.31	0.50	-	0.18	0.21	0.50
F	-	-	-	-	0.02	-	-	-	-	-	-	-	-	-	-	-	-	-	0.18	-	-
G	-	-	-	-	-	-	-	-	-	-	0.17	0.43	0.50	0.70	0.75	0.69	0.48	-	0.02	0.08	-
H	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	0.08	-
<b>Pep-2</b>																					
(N)	(6)	(15)	(20)	(7)	(12)	(27)	(13)	(10)	(19)	(32)	(12)	(5)	(24)	(13)	(4)	(27)	(15)	(6)	(27)	(8)	(8)
A	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
<b>Pgd</b>																					
(N)	(13)	(15)	(20)	(10)	(23)	(26)	(22)	(10)	(19)	(32)	(12)	(7)	(24)	(20)	(4)	(34)	(21)	(6)	(31)	(12)	(10)
A	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.08	-	-	-
B	-	-	-	-	-	0.08	-	-	-	-	0.04	-	-	-	0.13	0.02	0.02	0.17	-	-	-
C	0.23	-	0.02	-	0.02	0.23	0.07	0.20	0.05	0.09	0.96	0.93	0.96	0.95	0.87	0.85	0.86	0.58	0.21	-	-
D	0.77	1.00	0.98	1.00	0.98	0.69	0.93	0.80	0.95	0.91	-	0.07	0.04	0.05	-	0.13	0.12	0.17	0.79	-	0.05
E	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	1.00	0.95
<b>Pgi</b>																					
(N)	(14)	(15)	(20)	(10)	(23)	(26)	(21)	(10)	(19)	(32)	(12)	(7)	(23)	(20)	(4)	(34)	(21)	(6)	(31)	(12)	(10)
A	-	-	-	-	-	0.02	0.03	-	-	-	-	-	-	-	-	-	-	-	-	-	-
B	0.28	0.10	0.07	0.10	0.13	0.13	0.14	0.15	0.05	0.08	-	-	-	-	-	-	-	-	-	-	-
C	-	-	-	-	-	0.13	-	-	0.03	0.12	-	-	-	-	0.13	-	-	-	-	0.08	-
D	0.68	0.87	0.88	0.90	0.78	0.64	0.83	0.85	0.89	0.75	-	-	0.02	0.10	-	0.04	0.07	0.92	0.18	-	-
E	0.04	0.03	0.05	-	0.09	0.06	-	-	0.03	0.05	1.00	1.00	0.96	0.87	0.87	0.96	0.93	0.08	0.11	0.92	1.00
F	-	-	-	-	-	0.02	-	-	-	-	-	-	0.02	0.03	-	-	-	-	0.71	-	-
<b>Pgm</b>																					
(N)	(11)	(15)	(20)	(6)	(16)	(25)	(15)	(10)	(19)	(32)	(12)	(7)	(24)	(20)	(4)	(34)	(21)	(6)	(25)	(12)	(10)
A	0.09	0.03	-	-	-	0.02	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
B	-	0.07	-	0.08	0.06	0.08	0.07	-	-	0.09	-	-	-	-	-	-	-	-	-	-	-
C	0.14	0.30	0.20	0.25	0.25	0.36	0.10	0.20	0.16	0.39	-	-	-	-	-	-	-	-	-	-	-
D	0.32	0.23	0.55	0.67	0.47	0.32	0.27	0.10	0.39	0.46	-	-	-	-	-	-	-	-	-	-	-
E	0.45	0.37	0.20	-	0.22	0.16	0.56	0.70	0.32	0.06	-	-	-	-	-	-	-	-	-	-	-
F	-	-	0.05	-	-	0.06	-	-	0.13	-	-	-	-	-	-	-	-	0.25	-	1.00	1.00
G	-	-	-	-	-	-	-	-	-	-	0.88	1.00	0.96	1.00	0.87	0.91	0.88	0.75	0.02	-	-
H	-	-	-	-	-	-	-	-	-	-	0.12	-	0.04	-	0.13	0.09	0.12	-	0.06	-	-
I	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.92	-	-
<i>H<sub>o</sub></i>	0.23	0.16	0.15	0.17	0.19	0.24	0.18	0.16	0.21	0.23	0.11	0.11	0.13	0.11	0.10	0.10	0.14	0.11	0.20	0.16	0.14
<i>H<sub>e</sub></i>	0.26	0.17	0.16	0.17	0.20	0.28	0.21	0.18	0.24	0.24	0.11	0.14	0.11	0.10	0.09	0.11	0.14	0.13	0.21	0.15	0.11

0.05,  $df = 2.41$ ,  $p > 0.90$ ; *C. gigas*:  $F_{IS} = 0.04$ ,  $\chi^2 = 0.12$ ,  $df = 3.32$ ,  $p > 0.90$ ; *Ostrea* sp.:  $F_{IS} = -0.22$ ,  $\chi^2 = 0.47$ ,  $df = 0.32$ ,  $p > 0.40$ ).

We found 3 diagnostic loci that distinguished *Crassostrea rhizophorae* from *C. virginica*, 5 that distinguished *C. rhizophorae* from *C. brasiliana*, 6 that distinguished *C. brasiliana* from *C. virginica*, and 6 that distinguished *C. virginica* from *C. gigas* (Table 3). Although specimens of *C. virginica* and *C. brasiliana* are morphologically very similar, the former seems to be genetically more closely related to *C. rhizophorae* (Fig. 3). *C. brasiliana* and *C. paraibanensis* were genetically indistinguishable ( $D_{Nei} = 0$ ).

Intraspecific genetic distances were short among Brazilian populations of *Crassostrea brasiliana* ( $D_{Nei} = 0$  to 0.03) and of *C. rhizophorae* ( $D_{Nei} = 0$  to 0.08) (Fig. 3). Despite the large geographic distances involved (up to 9000 km), the genetic distances separating *C. rhizophorae* populations from Panama and Brazil were also small ( $D_{Nei} = 0.02$  to 0.08).

Populations of both species were found to be genetically structured in the Western Atlantic. Eight genetically different groups could be detected in *Crassostrea rhizophorae*: (1) Panama; (2) RN1; (3) RN2; (4) PB1, PB2; (5) BA2; (6) RJ1, SP1; (7) PR1; and (8) PR2 (SAMOVA:  $F_{CT} = 0.09$ ;  $p < 0.001$ ). Three genetic stocks were observed in *C. brasiliana*: (1) PA; (2) PB1, PE, BA1; and (3) SP2, PR1 (SAMOVA:  $F_{CT} = 0.08$ ,  $p < 0.01$ ).


Mantel tests for both species showed significant isolation by distance (IBD) relationships (*C. rhizophorae*:  $r = 0.53$ ,  $p < 0.05$ ; *C. brasiliana*:  $r = 0.77$ ,  $p < 0.01$ ; Fig. 4), even after excluding the very divergent Panamanian *C. rhizophorae* population ( $r = 0.36$ ,  $p < 0.05$ ).

### DNA markers

Analyses with the 3 DNA markers showed congruent results regarding interspecific relationships. ITS-2 intraspecific variation was low in all species (*Crassostrea rhizophorae*: 4 haplotypes,  $h = 0.271$ ,  $\pi = 0.001$ ; *C. virginica*: 1 haplotype; *C. brasiliana*/*C. paraibanensis*: 1 haplotype; and *C. gasar*: 2 haplotypes,  $h = 0.400$ ,  $\pi = 0.003$ ). *C. brasiliana*, *C. paraibanensis* and *C. gasar* were genetically extremely similar ( $K2P < 0.008$ ), whereas the other species could be readily distinguished by their ITS-2 sequences (Fig. 5).

We found 28 different COI haplotypes in the Western Atlantic populations of *Crassostrea rhizophorae* ( $h = 0.866$ ,  $\pi = 0.007$ ), 10 haplotypes in the South American populations of *C. brasiliana* ( $h = 0.590$ ,  $\pi = 0.003$ ), 3 haplotypes in the Brazilian population of *C. paraibanensis* ( $h = 0.524$ ,  $\pi = 0.001$ ), and only 1 haplotype in the *C. gasar* population from Joal-Fadiouth (Senegal, Africa). As observed with the ITS-2 sequences, *C. brasiliana*, *C. paraibanensis* and *C. gasar* were very

Fig. 3. *Crassostrea* spp. and *Ostrea* sp. UPGMA tree based on genetic distances (Nei 1978) between *Crassostrea* and *Ostrea* populations








Fig. 4. *Crassostrea* spp. Pairwise comparison between genetic (linearized  $F_{ST}/\Phi_{ST}$ ) and geographical (linearized km) distances based on nuclear (allozymes) and mitochondrial (COI) markers

similar in their COI sequences ( $K2P < 0.008$ ). Again, COI sequences clearly differentiated the other analyzed *Crassostrea* species ( $K2P$  varied between 0.16 and 0.27; Fig. 6). Intraspecific pairwise divergences of *C. rhizophorae* were low both for the Brazilian ( $K2P < 0.008$ ) and the Caribbean (Panama,  $K2P < 0.003$ ) populations. In contrast, a large differentiation ( $K2P = 0.018$  to 0.025) was found between sequences from Brazilian and Panamanian specimens, which clustered in separate clades with high bootstrap support in the neighbor-joining and maximum likelihood trees (Fig. 6).

With the exception of *Crassostrea brasiliana/C. paribanensis/C. gasar*, which had identical 16S sequences, the other *Crassostrea* species were well differentiated (Fig. 7). Interspecific sequence differentiation was 0.036 between *C. rhizophorae* and *C. virginica* and 0.118 between *C. rhizophorae* and *C. brasiliana*. No intraspecific variation was found in the 16S sequences analyzed.

Populations of both *Crassostrea rhizophorae* and *C. brasiliana* were found to be genetically structured in the Western Atlantic. Two highly divergent genetic




Fig. 5. *Crassostrea* spp. ITS-2-based (521 bp) tree. The numbers below the branches are bootstrap values (1000 replicates) for neighbor-joining and maximum likelihood trees, respectively

groups could be detected in *C. rhizophorae*: (1) Panama and (2) Brazil (SAMOVA COI data:  $F_{CT} = 0.894$ ,  $p < 0.0001$ ; Fig. 8). Excluding Panama from the analysis, 10 genetic stocks could be detected in *C. rhizophorae*: (1) CE; (2) RN1; (3) RN2; (4) PB1; (5) PB2; (6) PE; (7) BA2; (8) ES, RJ2, SP1; (9) RJ1; and (10) PR1, PR2 (SAMOVA COI data:  $F_{CT} = 0.156$ ,  $p < 0.001$ ). These population groupings are similar to those observed in the allozyme analyses. Three genetic stocks were observed in *C. brasiliana*: (1) PA, PB1, PE, BA1; (2) RJ3; and (3) SP2, PR1, PR2 (SAMOVA COI data:  $F_{CT} = 0.724$ ,  $p < 0.001$ ; Fig. 8). The *C. gasar* population from Joal-Fadiouth formed a fourth group when included in the analyses (SAMOVA COI data: *C. brasiliana/C. gasar*:  $F_{CT} = 0.738$ ,  $p < 0.001$ ; Fig. 8). As seen with allozymes, a significant correlation between geographic and genetic distances was observed in the COI analyses for *C. brasiliana* (Mantel test:  $r = 0.713$ ,  $p < 0.0001$ ; Fig. 4), whereas in *C. rhizophorae* the correlation was only sig-

nificant when including the highly differentiated Panamanian population (Fig. 4).

A significant departure from neutrality was observed for the *Crassostrea rhizophorae* populations from Panama (Tajima's  $D = -1.43$ ,  $p = 0.03$ ; Fu's  $F_S = -26.30$ ,  $p = 0$ ) and Brazil ( $D = -2.35$ ,  $p < 0.001$ ;  $F_S = -27.84$ ,  $p = 0$ ), and for *C. brasiliana* from northern and northeastern ( $D = -2.09$ ,  $p = 0.002$ ;  $F_S = -6.62$ ,  $p = 0.001$ ) and southeastern and southern ( $D = -1.67$ ,  $p = 0.026$ ;  $F_S = -0.60$ ,  $p = 0.209$ ) Brazilian populations. A unimodal distribution was observed in the mismatch analyses both for *C. rhizophorae* and *C. brasiliana*, closely matching the expected distributions under the sudden expansion model. Time estimates for population expansion were 27 thousand years ago (kya) for *C. rhizophorae* from Panama ( $\tau = 0.44$ ) and 107 kya for *C. rhizophorae* from Brazil ( $\tau = 1.72$ ). For *C. brasiliana*, estimates of time since expansion were 171 kya for the N/NE population ( $\tau = 2.75$ ) and 186 kya for the SE/S population ( $\tau = 3.00$ ).

## DISCUSSION

### Synonymy of *Crassostrea brasiliana*, *C. paraibanensis* and *C. gasar*

The 3 Western Atlantic *Crassostrea* species studied in this work, *C. rhizophorae*, *C. brasiliana* and *C. virginica*, could be readily distinguished by both nuclear (allozymes and ITS-2) and mitochondrial (COI and 16S) markers. On the other hand, *C. brasiliana* and the African *C. gasar* clustered together in all DNA analyses, which indicates that they are conspecific, as previously suggested (Verdon 2000, Lapègue et al. 2002) based on smaller data sets. Additionally, oysters found on rocks in the subtidal zone of the same shore where *C. paraibanensis* was described (Singarajah 1980) were genetically indistinguishable from *C. brasiliana/C. gasar*. We can safely conclude, therefore, that the nominal species *C. paraibanensis* and *C. brasiliana* are junior synonyms of *C. gasar*, which, henceforth, will be the name used to denominate both the South American and African oysters.

Moreau (2001) used restriction fragment length polymorphism (RFLP) analyses of ITS-2 to compare African and South American oysters identified as *Crassostrea gasar* and reported 2 different restriction patterns, H1 and H2, in the African specimens, but only H2 in the South American specimens. Because the African populations of *C. gasar* were more polymorphic, the author suggested that oysters from South America might have originated from African oysters.




Fig. 6. *Crassostrea* spp. Cytochrome *c* oxidase subunit I-based (648 bp) tree. The numbers on the branches are bootstrap values (1000 replicates) for neighbor-joining and maximum likelihood trees, respectively. 'a' and 'b' indicate the presence of haplotypes with individuals from more than one region (a: Southeast N = 2; b: Northeast N = 4)




Fig. 7. *Crassostrea* spp. 16S-based (495 bp) neighbor-joining tree. The numbers below the branches are bootstrap values (1000 replicates). The numbers in parentheses represent the total number of individuals sequenced with each haplotype

The high similarity between populations of the same oyster species on both sides of the South Atlantic indicates a recent dispersal event. This could have been either a naturally occurring event, such as the dispersal of larvae by the equatorial current (Lapègue et al. 2002) or of adults by rafting (Ó Foighil et al. 1999). Alternatively, it might be the result of the anthro-

pogenic transport of larvae in ballast water or of adults incrusting ship hulls as suggested by Lapègue et al. (2002).

The fossil records identified as *Crassostrea gasar* from Senegal (near Bas-soul) (Demarcq & Demarcq 1992) and as *C. 'brasiliiana'* from Brazil (S/SE regions) (Fairbridge 1976), date from the Holocene period (4000 to 6000 yr ago), thus refuting the anthropogenic transport hypothesis for the occurrence of *C. gasar* on both continents. In any case, the very high genetic similarity, across nuclear and mitochondrial genes, rules out vicariance as the origin of the African and South American oyster populations, since the 2 continents have been separated for over 16 million yr.

Cytochrome oxidase gene sequences from African *Crassostrea gasar* specimens were more similar to those from *C. gasar* from N/NE Brazil than to sequences from other parts of the Brazilian coast. This is compatible with colonization by natural dispersal from Africa through the westerly flowing Equatorial Current, which arrives in Brazil on its northeastern



### Phylogenetic relationships among Atlantic *Crassostrea* species and fossil records

All phylogenetic analyses demonstrated that the morphologically highly similar species, *Crassostrea rhizophorae* and *C. virginica*, are sister species genetically very different from *C. gasar*. However, *C. gasar* still groups with the other Atlantic oysters. The small genetic distances (allozymes:  $D_{Nei} = 0.02$  to 0.08) found between *C. rhizophorae* populations from Panama and Brazil (6000 to 9000 km apart) are similar to the genetic distances observed in comparisons of other marine organisms in those regions (Lazoski et al. 2001, Williams et al. 2001, Gusmão et al. 2006). This similarity provides additional support for the hypothesized close relationship of Caribbean and Brazilian faunas (Briggs 1974).

*Crassostrea* oysters, like other ostreids, are believed to have originated during the Late Cretaceous, like e.g. *C. soleniscus* (Meek, 1871) and *C. cusseta* Sohl & Kauffman, 1964 (Stenzel 1971), or earlier, in the Mid Jurassic period, like *C. tetoriensis* Komatsu & Chinzei, 2002 (Komatsu et al. 2002). During the Oligocene and Miocene periods, there was a great variety of oyster species, most of which had large and thick shells and inhabited littoral areas in the Western Atlantic (Kirby 2001). Some of these oysters probably gave rise to the Atlantic *Crassostreini* lineage after the Tethys Seaway closure (Ó Foighil & Taylor 2000).

Fossil records from the eastern coast of the USA (from the Priaborian period, Late Eocene, 37 million years ago, Ma) show the presence of a very large oyster, *Crassostrea gigantissima* (Finch, 1824), which was replaced by oysters similar to *C. virginica* in the Early Miocene strata (Aquitanian period, 23 Ma; Sohl & Kauffman 1964, Lawrence 1995, Kirby 2001). As a result, *C. virginica* is usually seen as a direct descendant of *C. gigantissima* or of an extinct common ancestor (Sohl & Kauffman 1964, Lawrence 1995, Kirby 2000). The American fossils identified as *C. virginica*, both from the Miocene and from the Holocene periods, had thin shells, whereas the Pliocene and Pleistocene '*C. aff. virginica*' specimens from the USA and Caribbean are thick-shelled (Kirby 2000, 2001). If this is correct, it means that *C. virginica* changed back from thin to thick shells during the Miocene-Holocene period. However, this conjecture is not consistent with the reported tendency of large and thick-shelled oysters to evolve into smaller and thinner-shelled species (Kirby 2000). This raises doubt as to the presence of *C. virginica* in the Miocene strata.

The divergence times estimated through allozymes and mitochondrial data indicate that *Crassostrea virginica* and *C. rhizophorae* diverged in the Pliocene period (6.8 to 3.2 Ma), after the evolutionary split in the lineages leading to *C. gasar* and the ancestor of *C. virginica*

and *C. rhizophorae*, which happened between the Miocene and the early Pliocene periods (9.5 to 4.6 Ma).

Based on the morphological and ecological similarities of *Crassostrea gasar* and *C. virginica* (Sandison & Hill 1966), it is important to consider the presence of *C. gasar* in the fossil record. This has not been done previously, because *C. gasar* was not considered a valid species in the Western Atlantic, so its morphological features (like the presence of thick and large shells) were only considered a part of the phenotypic space occupied by *C. virginica* and *C. rhizophorae*.

Pleistocene and Pliocene thick- and large-shelled oysters from the Atlantic and Caribbean (also identified as '*Crassostrea virginica*' and '*C. aff. virginica*'; Kirby 2000, 2001, Kirby & Jackson 2004) can be either *C. virginica* or *C. gasar*. This casts doubt on the identification of thick-shelled fossil oyster specimens as *C. virginica*, which means that the identification of fossil oysters, at least in the USA and Caribbean, should be revised.

Fossils identified as *Crassostrea virginica* date from the same period as those of *C. titan* (Conrad, 1853) from the state of California (USA) and *C. cahobasensis* (Pilsbry & Brown, 1917) from the Caribbean, which have large and thick shells (Kirby 2001) like those of *C. gasar*. Those shell features have been used to infer the habitat and other ecological characteristics of fossil oysters (Kirby 2000), and to aid in their classification (Sohl & Kauffman 1964, Stenzel 1971). However, the use of shell characteristics to distinguish *Crassostreina* genera has been criticized (Lawrence 1995). As the identification of extant oyster species based solely on their external shell features can clearly be misleading, correctly identifying and classifying fossil oysters based on those same shell characteristics is a difficult task (Lawrence 1995, Valentine et al. 2006).

### Geographical distribution, isolation by distance, and historical demography of Western Atlantic *Crassostrea rhizophorae* and *C. gasar*

*Crassostrea rhizophorae* was found in 13 of 21 locations studied, from the Caribbean (Panama) to southern Brazil (Paraná State, 25° 51' S; 48° 35' W), and *C. gasar* was found in 12 sites along the Brazilian coast ranging from the northern (Pará State, 01° 25' S; 48° 28' W) to the southern (Paraná State, 25° 51' S; 48° 35' W) regions. *C. rhizophorae* was only found in the intertidal environments. In contrast, *C. gasar*, which is known to live mainly in the subtidal zone, was also found in 8 intertidal sites, and in 2 of them (Paraíba State, northeastern Brazil, and Paranaguá Bay, southern Brazil) it was attached alongside *C. rhizophorae* on mangrove tree roots. Due to their large phenotypic plasticity, it is very difficult to correctly identify even adult specimens of

*C. gasar* and *C. rhizophorae* where they coexist. Hence, the confirmation, here, of the specific status of *C. rhizophorae* and *C. gasar*, the expansion of the known distribution of *C. gasar* to northern and northeastern Brazil, and the evidence of their occurrence in different habitats have important implications for biological and ecological studies of Latin American oysters. Due to their economic importance, the taxonomic separation of these oysters has consequences for biomonitoring and aquaculture programs (Ignacio et al. 2000, Rebelo et al. 2003).

A genetic study of allozymes from *Crassostrea rhizophorae* populations showed a weak genetic heterogeneity between populations near Rio de Janeiro (Southeast Brazil) and Paranaguá Bay (South Brazil) (Ignacio et al. 2000). Similar results were observed for other *Crassostrea* species (Buroker et al. 1979). However, the present analyses using a larger number of populations have unveiled significant intraspecific genetic differences both within *C. rhizophorae* and within *C. gasar*. Almost the same genetic partitioning detected by allozyme analyses of *C. rhizophorae* (one group in Panama and various in Brazil) and of *C. gasar* (at least one group each in northern/northeastern and south/southeastern Brazil) were also found by COI sequencing analyses.

There are many factors that can influence genetic population structure in marine organisms, including limited dispersal capability and the existence of extrinsic barriers to gene flow, such as temperature, salinity, and ocean currents (Launey et al. 2002), as well as historical factors, such as climate and marine current oscillations, and sea level changes (Rocha et al. 2008). Sampling strategies and genetic patchiness must also be taken into account (Arnaud-Haond et al. 2008).

Some intraspecific genetic differences have been observed in *Crassostrea rhizophorae* from the Caribbean (Hedgecock & Okazaki 1984) and between Atlantic and Gulf of Mexico *C. virginica* (Hare & Avise 1998, Varney et al. 2009). As the larvae of *Crassostrea* are planktonic and have great dispersal capability (for instance, *C. virginica* larvae can spend 2 to 3 wk in the plankton; Kennedy 1996), the high levels of genetic differentiation found in these works and in the present study are surprising.

The isolation by distance observed here for *Crassostrea rhizophorae* and *C. gasar* indicates that restricted larval dispersal may be one of the factors responsible for the genetic structuring of these species. Isolation by distance has also been detected in other oyster species, such as *Ostrea edulis* (Launey et al. 2002), *C. virginica* (Rose et al. 2006), and *C. ariakensis* (Xiao et al. 2010).

Mismatch distribution analyses suggest that the population history of both species is characterized by expansion events, which are congruent with changes

(ocean currents and sea levels) in the Late Pleistocene. Unfortunately, information about the biology of South American oyster species is still very scarce, which makes discussing the possible causes of the observed population heterogeneity difficult. Nevertheless, despite the factors responsible for the observed structuring patterns, our results show that those genetically distinct *Crassostrea rhizophorae* and *C. gasar* groups should be considered as discrete management units and this should be taken into account for future aquaculture programs and fisheries management of both resources, especially considering that these species usually display different growth rates and tolerances for temperature and salinity (Absher 1989, Melo et al. 2010b).

*Acknowledgements.* We thank T. Absher for invaluable discussions on oyster systematics and C. Melo for help in the identification of oysters, as well as H. Cunha for helpful suggestions that improved an earlier version of the manuscript and F. Monteiro for providing sequencing support. We also thank A. Freire, A. Pereira, A. Puchnick Legat, B. Ignacio, C. Tureck, C. Zilberberg, D. Almeida, F. Nunes, J. Legat, M. Klautau, M. Miranda, M. Oliveira, M. Rebelo, N. Knowlton, P. Paiva, P. Vianna, S. Christo, and T. Absher for help with sample collection. This work is part of the PhD thesis of the first author at the Genetics Department of the Federal University of Rio de Janeiro (UFRJ) and was supported by grants from the Brazilian CAPES, CNPq, FAPERJ, FINEP, FUJB, MCT (CT-AGRO/CT-HIDRO), MPA, and PADCT.

#### LITERATURE CITED

- Absher TM (1989) Populações naturais de ostras do gênero *Crassostrea* do litoral do Paraná—desenvolvimento larval, recrutamento e crescimento. PhD thesis, University of São Paulo
- Arnaud-Haond S, Vonau V, Rouxel C, Bonhomme F, Prou J, Goyard E, Boudry P (2008) Genetic structure at different spatial scales in the pearl oyster (*Pinctada margaritifera cumingi*) in French Polynesian lagoons: beware of sampling strategy and genetic patchiness. *Mar Biol* 155: 147–157
- Batista FM, Leitão A, Huvet A, Lapègue S, Heurtebise S, Boudry P (2006) The taxonomic status and origin of the Portuguese oyster *Crassostrea angulata* (Lamarck, 1819). *Oyster Res Inst News* 18:3–10
- Belkhir K, Borsa P, Chikhi L, Raufaste N, Bonhomme F (2002) GENETIX 4.04, logiciel sous Windows pour la génétique des populations. Montpellier
- Boudry P, Heurtebise S, Collet B, Cornette F, Gerard A (1998) Differentiation between populations of the Portuguese oyster, *Crassostrea angulata* (Lamarck) and the Pacific oyster, *Crassostrea gigas* (Thünberg), revealed by mtDNA RFLP analysis. *J Exp Mar Biol Ecol* 226:279–291
- Boudry P, Heurtebise S, Lapègue S (2003) Mitochondrial and nuclear DNA sequence variation of presumed *Crassostrea gigas* and *C. angulata* specimens: a new oyster species in Hong Kong? *Aquaculture* 228:15–25
- Briggs JC (1974) *Marine zoogeography*. McGraw-Hill, New York
- Buroker NE, Hershberger WK, Chew KK (1979) Population genetics of the family Ostreidae. II. Interspecific studies

- of the genera *Crassostrea* and *Saccostrea*. *Mar Biol* 54: 171–184
- Carranza A, Defeo O, Beck M (2009) Diversity, conservation status and threats to native oysters (Ostreidae) around the Atlantic and Caribbean coasts of South America. *Aquat Conserv: Mar Freshw Ecosyst* 19:344–353
- Carriker MR, Gaffney PM (1996) A catalogue of selected species of living oysters (Ostreacea) of the world. In: Kennedy VS, Newell RIE, Eble AF (eds) *The eastern oyster: C. virginica*. Maryland Sea Grant College, Maryland, p 1–18
- Cenik C, Wakeley J (2010) Pacific salmon and the coalescent effective population size. *PLoS ONE* 5:e13019
- Clement M, Posada D, Crandall KA (2000) TCS: a computer program to estimate gene genealogies. *Mol Ecol* 9: 1657–1659
- Demarcq H, Demarcq G (1992) The *Crassostrea gasar* (Bivalvia) biostrome of the Holocene from Sine-Saloum (Senegal)—new data and ecostratigraphical interpretation. *Geobios* 25:225–250
- Dixon DR, Solé-Cava AM, Pascoe PL, Holland PWH (1995) Periostracal adventitious hairs on spat of the mussel *Mytilus edulis*. *J Mar Biol Assoc UK* 75:363–372
- Dupanloup I, Schneider S, Excoffier L (2002) A simulated annealing approach to define the genetic structure of populations. *Mol Ecol* 11:2571–2581
- Eldon B, Wakeley J (2006) Coalescent processes when the distribution of offspring number among individuals is highly skewed. *Genetics* 172:2621–2633
- Excoffier L, Laval G, Schneider S (2005) Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evol Bioinform Online* 1:47–50
- Fairbridge RW (1976) Shellfish-eating pre-ceramic Indians in coastal Brazil. *Science* 191:353–359
- FAO (2010) Fishstat Plus 2.30: Universal software for fishery statistical time series. Fisheries Department, Fishery Information, Data and Statistics Unit, Rome
- Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* 17:368–376
- Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R (1994) DNA primers for amplification of mitochondrial cytochrome *c* oxidase subunit I from diverse metazoan invertebrates. *Mol Mar Biol Biotechnol* 3:294–299
- Fu YX (1997) Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics* 147:915–925
- Gusmão J, Solé-Cava AM (2002) Um sistema de diagnóstico molecular para a identificação de espécies comerciais de camarões marinhos brasileiros. In: Blas I (ed) CIVA 2002 ([www.civa2002.org](http://www.civa2002.org)), Zaragoza, p 754–764
- Gusmão J, Lazoski C, Monteiro FA, Solé-Cava AM (2006) Cryptic species and population structuring of the Atlantic and Pacific seabob shrimp species, *Xiphopenaeus kroyeri* (Heller 1862) and *X. riveti* (Bouvier 1907). *Mar Biol* 149: 491–502
- Hare MP, Avise JC (1998) Population structure in the American oyster as inferred by nuclear gene genealogies. *Mol Biol Evol* 15:119–128
- Harry HW (1985) Synopsis of supraspecific classification of living oysters (Bivalvia: Gryphaeidae and Ostreidae). *Veliger* 28:121–158
- Hasegawa M, Kishino H, Yano T (1985) Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *J Mol Evol* 22:160–174
- Hedgecock D (1994) Does variance in reproductive success limit effective population sizes of marine organisms? In: Beaumont AR (ed) *Genetics and evolution of aquatic organisms*. Chapman & Hall, London, p 122–134
- Hedgecock D, Okazaki NB (1984) Genetic diversity within and between populations of American oysters (*Crassostrea*). *Malacologia* 25:535–549
- Hedgecock D, Launey S, Pudovkin AI, Naciri Y, Lapègue S, Bonhomme F (2007) Small effective number of parents ( $N_b$ ) inferred for a naturally spawned cohort of juvenile European flat oysters *Ostrea edulis*. *Mar Biol* 150:1173–1182
- Hoover CA, Gaffney PM (2005) Geographic variation in nuclear genes of the eastern oyster *Crassostrea virginica* Gmelin. *J Shellfish Res* 24:103–112
- Ignacio BL, Absher TM, Lazoski C, Solé-Cava AM (2000) Genetic evidence for the presence of two species of *Crassostrea* (Bivalvia: Ostreidae) on the coast of Brazil. *Mar Biol* 136:987–992
- Jensen JL, Bohonak AJ, Kelley ST (2005) Isolation by distance, web service. *BMC Genet* 6:13
- Johansson ML, Banks MA, Glunt KD, Hassel-Finnegan HM, Buonaccorsi VP (2008) Influence of habitat discontinuity, geographical distance, and oceanography on fine-scale population genetic structure of copper rockfish (*Sebastes caurinus*). *Mol Ecol* 17:3051–3061
- Kennedy VS (1996) The biology of larvae and spat. In: Kennedy VS, Newell RIE, Eble AF (eds) *The eastern oyster: Crassostrea virginica*. Maryland Sea Grant College, Maryland, p 371–422
- Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16:111–120
- Kirby MX (2000) Paleoeological differences between Tertiary and Quaternary *Crassostrea* oysters, as revealed by stable isotope sclerochronology. *Palaios* 15:132–141
- Kirby MX (2001) Differences in growth rate and environment between Tertiary and Quaternary *Crassostrea* oysters. *Paleobiology* 27:84–103
- Kirby MX, Jackson JBC (2004) Extinction of a fast-growing oyster and changing ocean circulation in Pliocene tropical America. *Geology* 32:1025–1028
- Komatsu T, Chinzei K, Zakhara MS, Matsuoka H (2002) Jurassic soft-bottom oyster *Crassostrea* from Japan. *Palaeontology* 45:1037–1048
- Lapègue S, Boutet I, Leitão A, Heurtebise S, Garcia P, Thiriot-Quévieux C, Boudry P (2002) Trans-Atlantic distribution of a mangrove oyster species revealed by 16S mtDNA and karyological analyses. *Biol Bull* 202:232–242
- Launey S, Ledu C, Boudry P, Bonhomme F, Naciri-Graven Y (2002) Geographic structure in the European flat oyster (*Ostrea edulis* L.) as revealed by microsatellite polymorphism. *J Hered* 93:331–338
- Lawrence DR (1995) Diagnosis of the genus *Crassostrea* (Bivalvia, Ostreidae). *Malacologia* 36:185–202
- Lazoski C (2004) Sistemática molecular e genética populacional de ostras brasileiras (*Crassostrea* spp.). PhD thesis, Federal University of Rio de Janeiro
- Lazoski C, Solé-Cava AM, Boury-Esnault N, Klautau M, Russo CAM (2001) Cryptic speciation in a high gene flow scenario in the oviparous marine sponge *Chondrosia reniformis*. *Mar Biol* 139:421–429
- Leitão A, Chaves R, Santos S, Guedes-Pinto H, Boudry P (2007) Interspecific hybridization in oysters: restriction enzyme digestion chromosome banding confirms *Crassostrea angulata* × *C. gigas* F1 hybrids. *J Exp Mar Biol Ecol* 343:253–260
- Littlewood DTJ (1994) Molecular phylogenetics of cupped oysters based on partial 28S rRNA gene sequences. *Mol Phyl Evol* 3:221–229
- Manchenko GP (1994) *A handbook of detection of enzymes on electrophoretic gels*. CRC Press, London
- Marko PB (2002) Fossil calibration of molecular clocks and the divergence times of geminate species pairs separated by

- the Isthmus of Panama. *Mol Biol Evol* 19:2005–2021
- Melo AGC, Varela ES, Beasley CB, Schneider H and others (2010a) Molecular identification, phylogeny and geographic distribution of Brazilian mangrove oysters (*Crassostrea*). *Genet Mol Biol* 33:564–572
- Melo CMR, Solé-Cava AM, Lazoski C (2010b) *Crassostrea gigas* in natural oyster banks in southern Brazil. *Biol Invasions* 12:441–449
- Milbury CA, Gaffney PM (2005) Complete mitochondrial DNA sequence of the eastern oyster *Crassostrea virginica*. *Mar Biotechnol* 7:697–712
- Moreau D (2001) Etude de génétique des populations sur l'huître creuse de mangrove, *Crassostrea gasar* (Adanson, 1757), par l'apport du marqueur ITS2. Diplôme d'Etudes approfondies en Biologie des Populations et écosystèmes. Université de Sciences la Rochelle, France
- Nei M (1978) Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics* 89:583–590
- Nei M (1987) *Molecular evolutionary genetics*. Columbia University Press, New York
- Nielsen R, Wakeley J (2001) Distinguishing migration from isolation: a Markov chain Monte Carlo approach. *Genetics* 158:885–896
- Ó Foighil D, Marshall BA, Hilbish TJ, Pino MA (1999) Trans-Pacific range extension by rafting is inferred for the flat oyster *Ostrea chilensis*. *Biol Bull* 196:122–126
- Ó Foighil D, Taylor DJ (2000) Evolution of parental care and ovulation behavior in oysters. *Mol Phyl Evol* 15:301–313
- Palumbi SR (1996) Nucleic acids II: The polymerase chain reaction. In: Hillis DM, Moritz C (eds) *Molecular systematics*. Sinauer Associates, Massachusetts, p 205–247
- Palumbi SR (2003) Population genetics, demographic connectivity, and the design of marine reserves. *Ecol Appl* 13: S146–S158
- Pie MR, Ribeiro RO, Boeger WA, Ostrensky A, Falleiros RM, Angelo L (2006) A simple PCR-RFLP method for the discrimination of native and introduced oyster species (*Crassostrea brasiliiana*, *C. rhizophorae* and *C. gigas*; Bivalvia: Ostreidae) cultured in Southern Brazil. *Aquacult Res* 37: 1598–1600
- Posada D, Crandall KA (1998) MODELTEST: testing the model of DNA. *Bioinformatics* 14:817–818
- Rebelo MF, Pfeiffer WC, da Silva H, Moraes MO (2003) Cloning and detection of metallothionein mRNA by RT-PCR in mangrove oysters (*C. rhizophorae*). *Aquat Toxicol* 64:359–362
- Reece KS, Cordes JF, Stubbs JB, Hudson KL, Francis EA (2008) Molecular phylogenies help resolve taxonomic confusion with Asian *Crassostrea* oyster species. *Mar Biol* 153: 709–721
- Rios EC (1994) *Seashells of Brazil*. FURG, Rio Grande
- Rocha LA, Rocha CR, Robertson DR, Bowen BW (2008) Comparative phylogeography of Atlantic reef fishes indicates both origin and accumulation of diversity in the Caribbean. *BMC Evol Biol* 8:157
- Rogers AR, Harpending HC (1992) Population growth makes waves in the distribution of pairwise genetic differences. *Mol Biol Evol* 9:552–569
- Rose CG, Paynter KT, Hare MP (2006) Isolation by distance in the eastern oyster, *Crassostrea virginica*, in Chesapeake Bay. *J Hered* 97:158–170
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Sandison EE, Hill MB (1966) The distribution of *Balanus pallidus stutsburi* Darwin, *Gryphaea gasar* ((Adanson) Dautzenberg), *Mercierella enigmatica* Fauvel and *Hydroides uncinata* (Philippi) in relation to salinity in Lagos Harbour and adjacent creeks. *J Anim Ecol* 35:235–250
- Singarajah KV (1980) On the taxonomy, ecology and physiology of a giant oyster, *Crassostrea paraibanensis*, new species. *Bull Mar Sci* 30:833–841
- Sohl NF, Kauffman EG (1964) Giant upper Cretaceous oysters from the Gulf coast and Caribbean. *US Geol Surv Prof Pap* 483:1–22
- Stenzel HB (1971) Bivalvia Mollusca: oysters. In: Moore RC (ed) *Treatise on invertebrate paleontology*. GSA-University of Kansas Press, Lawrence, p 953–1217
- Swofford DL (1998) PAUP. Phylogenetic analysis using parsimony, version 4. Sinauer Associates, Sunderland, MA
- Swofford DL, Selander RB (1981) BIOSYS-1, a FORTRAN programme for the comprehensive analysis of electrophoretic data in population genetics and systematics. *J Hered* 72: 281–283
- Tajima F (1989) Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123:585–595
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24:1596–1599
- Templeton AR, Crandall KA, Sing CF (1992) A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping and DNA-sequence data. 3. Cladogram estimation. *Genetics* 132: 619–633
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The ClustalX Windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25:4876–4882
- Valentine JW, Jablonski D, Kidwell S, Roy K (2006) Assessing the fidelity of the fossil record by using marine bivalves. *Proc Natl Acad Sci USA* 103:6599–6604
- Varela ES, Beasley CR, Schneider H, Sampaio I, Marques-Silva NS, Tagliaro CH (2007) Molecular phylogeny of mangrove oysters (*Crassostrea*) from Brazil. *J Mollusc Stud* 73:229–234
- Varney RL, Galindo-Sanchez CE, Cruz P, Gaffney PM (2009) Population genetics of the eastern oyster *Crassostrea virginica* (Gmelin, 1791) in the Gulf of Mexico. *J Shellfish Res* 28:855–864
- Verdon B (2000) Phylogéographie des huîtres creuses des mangroves de l'Atlantique Sud par l'apport des marqueurs moléculaires. Diplôme d'Etudes Approfondies en Biologie des Populations et Ecosystèmes. Université de Sciences La Rochelle, France
- Waples RS (1987) A multispecies approach to the analysis of gene flow in marine shore fishes. *Evolution* 41:385–400
- Williams ST, Knowlton N, Weigt LA, Jara JA (2001) Evidence for three major clades within the snapping shrimp genus *Alpheus* inferred from nuclear and mitochondrial gene sequence data. *Mol Phyl Evol* 20:375–389
- Wright S (1978) *Evolution and the genetics of populations*, Vol 4. The University of Chicago Press, London
- Wu X, Xu X, Yu Z, Wei Z, Xia J (2010) Comparison of seven *Crassostrea* mitogenomes and phylogenetic analyses. *Mol Phyl Evol* 57:448–454
- Xiao J, Cordes JF, Wang H, Guo X, Reece KS (2010) Population genetics of *Crassostrea ariakensis* in Asia inferred from microsatellite markers. *Mar Biol* 157:1767–1781